# Joint Crediting Mechanism Approved Methodology KH\_AM004 "Reducing deforestation and forest degradation through forest conservation in Cambodia"

# A. Title of the methodology

Reducing deforestation and forest degradation through forest conservation in Cambodia, version 1.10

### **B.** Terms and definitions

Terms	Definitions		
Cambodia's official forest	Cambodia's official forest reference level (National FRL)		
reference level (National FRL)	is defined as the most recent and available forest reference		
	level submitted to the UNFCCC secretariat by Cambodia		
	and completed technical assessment by the UNFCCC's		
	assessment team.		
Cambodia's official forest map	A land use and land cover map, endorsed as an official		
	map, named Forest Cover, by the Government of		
	Cambodia. The generation of additional landcover data are		
	expected every 2 years, e.g., Forest Cover 2018 and Forest		
	Cover 2020.		
Forest classes	Classes of forests adopted in the establishment of		
	Cambodia's official forest reference level.		
Transition probability	A probability that a forest class will be changed to another		
	forest class, or be converted, to non-forest, based on the		
	historical pace of transition. Transition probability is		
	between 0 to 1.		

# C. Summary of the methodology

Items	Summary
Project activities	The project activities include forest conservation activities such as forest management and community livelihood development, including

(emission reduction measures)	agricultural improvement, eco-tourism development, marketing of non- timber forest products and formation/strengthening of community groups that lead to the reduction of emissions from deforestation and degradation.
Establishment of project reference level	<ul> <li>Two options have been identified to establish the project reference level, using data from Cambodia's official forest reference level (National FRL): <ul> <li>(1) applying the National FRL transition probabilities from forest to non-forest classes only, to the project area, (e.g., transitions <i>between</i> different forest classes are not included); and</li> <li>(2) applying all the National FRL transition probabilities among classes that would result in emissions to the project area.</li> </ul> </li> <li>Note that Option (1) accounts for emissions from only deforestation, while Option (2) accounts for emissions from forest degradation in form of transitions from one forest class to another, in addition to those from deforestation.</li> <li>Option 2 is similar to the method used in the Cambodia's National FRL submission in 2017, except that the National FRL includes removals as well; this methodology targets <i>only</i> emissions.</li> </ul>
Calculation of project net emissions	Project net emissions are estimated based on the result of monitoring of forest to non-forest classes (Option 1) and monitoring of forest to non- forest or another forest class (Option 2) within the project area, multiplied by the emission factors (including both above and below ground biomass) plus any displaced emissions within the displacement belt and GHG emissions due to the project activities inside the project area and the activity area in line with Joint Crediting Mechanism Guidelines for Developing Proposed Methodology for Reducing Emissions from Deforestation and Forest Degradation, and the Role of Conservation, Sustainable Management of Forests and Enhancement of Forest Carbon Stocks in Developing Countries (REDD-plus), (methodology guidelines). Displaced emissions are calculated as increases of emissions in the displacement belt compared to the reference emissions. Use of fuel and fertilizers, for the implementation of the project activities, are included as GHG emissions sources, and will be monitored and accounted as project emissions.
Monitoring parameters and methods	<ul> <li>Area converted from forest to non-forest (Option 1) or forest to another forest class (Option 2) in the project area</li> <li>Area converted from forest to non-forest (Option 1) or forest to another forest class (Option 2) in the displacement belt</li> <li>Quantity of fuel consumed in vehicle and/or equipment</li> <li>Number of vehicle and/or equipment</li> <li>Total travel distance of vehicle and/or total use hours of equipment</li> <li>Average specific energy consumption of vehicle and/or equipment</li> <li>Mass of synthetic fertilizer applied</li> <li>Mass of organic fertilizer applied and made from materials sourced from outside of the project area and the activity area</li> <li>Harvested annual dry matter yield of N-fixing crop per unit area, introduced by the project</li> <li>Total annual area harvested of N-fixing crop introduced by the project</li> <li>Ratio of above-ground residues to harvested yield for N-fixing crop</li> <li>Mass of calcic limestone, dolomite and urea fertilizer applied</li> </ul>

ļ

	<ul> <li>Nitrogen content of synthetic and organic fertilizer applied</li> <li>Nitrogen content of above-ground and below-ground residues for N-fixing crop</li> </ul>
Calculation of project emission reductions to be credited	A default discount factor of 20%, as defined in the methodology guidelines, is applied to project emissions reductions to account for the risk of reversal.

# **D.** Eligibility criteria

1

This methodology is applicable to projects that satisfy all of the following criteria.

Criterion 1	The project is to reduce deforestation and forest degradation through project		
	activities including forest management and community livelihood		
	development.		
Criterion 2	Cambodia's official forest reference (emission) level has been submitted to		
	UNFCCC, completed technical assessment by UNFCCC, and is publicly		
	available.		
Criterion 3	Cambodia's official forest map for the project start year or less than or equal		
	to two years old is available for the project participant.		
Criterion 4	Project activities do not include activities which lead to GHG emissions		
	within the project area and the project activity area, except for the use of fuel		
	or fertilizer including N-fixing crops.		

## E. Geographical Boundaries

Essential			
Geographical boundary	Requirements		
Project area	No requirements in addition to those described in paragraphs 17 to 22 of the methodology guidelines ver01.0 (JCM_KH_GL_PM_REDD+_ver01.0)		
Reference area	The land area of Cambodia is used to keep consistency with National FRL.		

#### Optional boundaries

Geographical boundary	Required (Y/N/TBD)	Additional requirements	
Activity area	Y	No requirements in addition to those described in paragraph 17 and 21 of the methodology guidelines ver01.0.	

Displacement helt	Y	Displacement belt is the forest which is located
Displacement beit		outside of the project area, where deforestation
		and forest degradation could occur due to the
		displacement of project activities, such as small-
		scale logging, agricultural encroachment, and
		collection of non-timber forest products. The
		displacement belt is delineated on the basis of
		information on the impact of project activities
		which are obtained from local experts and other
		sources. If other REDD+ projects exist within
		the belt, project areas of these projects are
		excluded from the displacement belt. If
		development plans, including Economic Land
		Concessions (ELCs), exist within the belt, those
		areas are also excluded because forest clearance
		is likely planned inside these areas regardless
		the project activities.

TBD: to be decided by the project participant

## F. Carbon pools and GHG sources

The net emission sources to be considered include all the following Carbon pools and GHG sources.

Project reference level			
Carbon pools and GHG sources		Included	Explanation
		(Y/N)	Explanation
	Above ground biomass	Y	This pool is expected to contribute significantly to emissions and emission reductions and is therefore included.
Carbon pools	Below ground biomass	Y	This pool is expected to contribute significantly to emissions and emission reductions and is therefore included.
	Dead wood	N	It is expected that this pool would have decreased in the absence of the project and, therefore, it is conservatively excluded.
	Litter	N	It is expected that this pool would have decreased in the absence of the project and, therefore, it is conservatively excluded.
	Soil organic carbon	N	It is expected that this pool would have decreased in the absence of the project and, therefore, it is conservatively excluded.
GHG			
sources			
Project net emissions			

Carbon pools and CUC sources		Included	Employedian
Carbor	i pools and GHG sources	(Y/N)	Explanation
Above gro Below gro Carbon pools Litter Soil organi	Above ground biomass	Y	This pool is expected to contribute significantly to emissions and emission reductions and is therefore included.
	Below ground biomass	Y	This pool is expected to contribute significantly to emissions and emission reductions and is therefore included.
	Dead wood	N	It is expected that this pool would have decreased in the absence of the project and, therefore, it is conservatively excluded.
	Litter	N	It is expected that this pool would have decreased in the absence of the project and, therefore, it is conservatively excluded.
	Soil organic carbon	N	It is expected that this pool would have decreased in the absence of the project, and therefore, it is conservatively excluded.
GHG	CO <sub>2</sub> emissions from combustion of fossil fuels		This GHG source is estimated if the project activities which include combustion of fossil fuels are implemented
sources	N <sub>2</sub> O and CO <sub>2</sub> emissions from fertilizer application		This GHG source is estimated if the project activities include fertilizer application.

### G. Establishment of project reference level

### G.1. Establishment of project reference level

<u>Reference area and reference period</u> are the same as the National FRL, and therefore the reference area is the total of Cambodia's land area. The reference period is the same as the reference period submitted in the National FRL to the UNFCCC. In the National FRL submitted in 2017; the reference period was 2006 to 2014.

<u>Calculation of transition probability</u> is done by applying one of two options provided in this methodology: (1) Option 1: using the National FRL transition probabilities from forest to non-forest classes only (i.e., transitions *between* different forest classes are not included); or (2) Option 2: using all the National FRL transition probabilities among classes that would result in

GHG emissions. The option selected should also be used for calculation of project net emissions in section H and for determining data and parameters fixed *ex ante* in section J. The option is selected in coordination with Government of Cambodia and project participants.

<u>Yearly reference emission</u> at the project reference level is calculated by applying the transition probabilities to areas of forests at the previous year and multiplying by emission factors.

To ensure the conservativeness of the calculation of the project emission reductions, dead wood, litter and soil organic carbon pools, of which carbon stocks would have decreased in the absence of the project, were excluded.

#### G.2. Calculation of project reference level

The project reference level in year y during the monitoring period is calculated as follows:  $RL_y = \Delta CS_{refy} * 44/12$  Equation 1

Where:

$RL_y$	Project reference level in year y; tCO <sub>2</sub>
$\Delta CS_{refy}$	Projected carbon stock change in the project area in year y; tC
44/12	Conversion factor of molecular weight of carbon to CO <sub>2</sub> ; dimensionless

 $\Delta CS_{ref y}$  is calculated by applying either Option 1 or Option 2 selected in coordination with government of Cambodia and project participants.

Option 1: Use the National FRL transition probabilities from forest to non-forest classes only

Carbon stock change in the project area in year *y* is calculated as follows:

For the project start year:

 $\Delta CS_{ref y} = \sum_{i} (A_{i 0} * P_{i} * d_{y} / d_{0 y} * EF_{i})$ Equation 2 For subsequent years to the project start year:

$$\Delta CS_{ref y} = \sum_{i} (A_{i y-1} * P_i * d_y / d_{0 y} * EF_i)$$
Equation 332

Where:

 $\Delta CS_{ref y}$ Projected carbon stock change in the project area in year y; tC $A_{i 0}$ Area of forest class i in the project area at the inception of the project; ha

$A_{iy\text{-}1}$	Area of forest class <i>i</i> in the project area in year <i>y</i> - <i>1</i> ; ha
i	Forest class in the project area; dimensionless
Pi	Annual transition probability from forest class <i>i</i> to non-forest within the reference
	area; dimensionless, 0-1
<u>d</u> y	Number of operating days in year y, days
<u>d</u> o y	Number of days in year y, days
EFi	Emission factor applicable for forest class $i$ ; tC ha <sup>-1</sup>

Area of forest class *i* is assumed to decrease every year due to deforestation, and therefore calculated as follows:

For the project start year:

 $A_{iy} = A_{iy-1} * (1 - P_i)$ 

 $A_{iy+1} = A_{i\neq 0} * (1 - P_i) \underline{d_y} / d_{0y}$ For subsequent years to the project start year:

Equation <u>44</u>3

Equation 5

Where:

A <sub>i y+1</sub>	_Area of forest class <i>i</i> in the project area in year <i>y</i> +1; ha
$A_{iy}$	Area of forest class <i>i</i> in the project area in year <i>y</i> ; ha
<u>A<sub>i 0</sub></u>	Area of forest class <i>i</i> in the project area at the inception of the project; ha
<u>A</u> <sub>i y-1</sub>	Area of forest class <i>i</i> in the project area in year <i>y</i> - <i>1</i> ; ha
$\mathbf{P}_{\mathrm{i}}$	Annual transition probability from forest class <i>i</i> to non-forest within the reference
	area; dimensionless, 0-1
<u>d</u> y	Number of operating days in year y, days
d <sub>o v</sub>	Number of days in year v, days

Please refer to section J for determination of  $EF_i$ ,  $A_{i0}$  (area at the inception of the project) and  $P_i$ .

Option 2: Use all the National FRL transition probabilities among classes that would result in GHG emissions.

To estimate the total emissions, the projected carbon stock change in year y in the project area is calculated as follows:

$$\Delta CS_{ref y} = \sum_{i} \sum_{j} cs_{ij y} \begin{cases} cs_{ij y}, & \text{if } cs_{ij y} > 0\\ 0, & \text{otherwise} \end{cases}$$
 Equation 64

Where:

 $\Delta CS_{refy}$  Projected carbon stock change in the project area in year y; tC

cs<sub>ij y</sub> Projected carbon stock change in the project area from changes of land use category *i* to *j* in year *y*; tC
i, j Land use category in the project area; dimensionless

The notation  $cs_{ijy}$  is an element of *i*-by-*j* cross-tabulation matrix  $MCS_y$  which is a product of element-wise multiplication (Hadamard product) of *i*-by-*j* matrices of area changes and emission factors.

$$MCS_{\nu} = MCA_{\nu} \cdot MEF$$

Equation 75

Where:

- MCS<sub>y</sub> i-by-j matrix in which each element is projected carbon stock change in the project area from changes of land use category i to j in the project area in year y,  $cs_{ijy}$ ; tC
- MCA<sub>y</sub> *i*-by-*j* matrix in which each element is projected area of land converted from land use category *i* to *j* in the project area in year *y*,  $ca_{ijy}$ ; ha
- MEF *i*-by-*j* matrix in which each element is emission factor for area of land converted from land use category *i* to *j*,  $EF_{ij}$ ; tC ha<sup>-1</sup>

Equation 5 is expressed as below:

CS <sub>11y</sub>	cs <sub>12y</sub>		cs <sub>1jy</sub>	ca <sub>11y</sub>	ca <sub>12y</sub>	 ca <sub>1jy</sub>	$\Gamma^{\text{EF}_{11}}$	EF <sub>12</sub>	 EF <sub>1j</sub> ]
cs <sub>21y</sub>	cs <sub>22y</sub>		cs <sub>2jy</sub>	ca <sub>21y</sub>	ca <sub>22y</sub>	 ca <sub>2jy</sub>	EF <sub>21</sub>	EF <sub>22</sub>	 EF <sub>2j</sub>
				-		 			 
cs <sub>i1y</sub>	cs <sub>i2y</sub>	•••	cs <sub>ijy</sub>	ca <sub>i1y</sub>	ca <sub>i2y</sub>	 ca <sub>ijy</sub>	EF <sub>i1</sub>	EF <sub>i2</sub>	 EF <sub>ij</sub> ]

For example, the  $cs_{IIy}$  is calculated as bellow:

$$cs_{11y} = ca_{11y} * EF_{11}$$

Projected area of land conversion in year y is calculated as the product between areas of land categories in the project area in year y-l and annual transition probabilities.

 $MCA_v = MA_{v-1}MP$ 

Equation 86

Where:

MCA <sub>y</sub>	<i>i</i> -by- <i>j</i> matrix in which each element is the projected area of land converted from
	land use category <i>i</i> to <i>j</i> in year <i>y</i> , $ca_{ijy}$ ; ha
$MA_{y-1}$	<i>i</i> -by- <i>j</i> diagonal matrix whose diagonal elements are areas of land use categories $i$
	in the project area in year <i>y</i> - <i>1</i> , A <sub>i y-1</sub> ; ha
MP	<i>i</i> -by- <i>j</i> matrix in which each element is the annual transition probability from land
	use category <i>i</i> to <i>j</i> , $p_{ij}$ ; dimensionless, 0-1

Equation 6 is expressed as below:

[ca <sub>11y</sub>	ca <sub>12y</sub>	 ca <sub>1jy</sub> ]		A <sub>1y-1</sub>	0	 0	<b>[</b> p <sub>11</sub>	p <sub>12</sub>	 p <sub>1j</sub> ]
ca <sub>21y</sub>	ca <sub>22y</sub>	 ca <sub>2jy</sub>	=	0	$A_{2y-1}$	 0	p <sub>21</sub>	p <sub>22</sub>	 p <sub>2j</sub>
		 				 		•••	 
ca <sub>i1y</sub>	ca <sub>i2y</sub>	 ca <sub>ijy</sub>		L O	0	 A <sub>iy-1</sub>	p <sub>i1</sub>	$\mathbf{p_{i2}}$	 p <sub>ij</sub>

For example, selected elements are calculated as bellow:

$$\begin{split} & ca_{11y} = A_{1y\text{-}1} \, * \, p_{11} + 0 \, * \, p_{21} + \ldots + 0 \, * \, p_{i1} \\ & ca_{21y} = 0 * \, p_{11} + A_{2y\text{-}1} \, * \, p_{21} + \ldots + 0 \, * \, p_{i1} \\ & ca_{22y} = 0 \, * \, p_{12} + A_{2y\text{-}1} \, * \, p_{22} + \ldots + 0 \, * \, p_{i2} \\ & ca_{ijy} = 0 \, * \, p_{1j} + 0 \, * \, p_{2j} + \ldots + A_{iy\text{-}1} \, * \, p_{ij} \end{split}$$

Area of each land use category is decreased due to transition to other land categories and increased due to transition from other land use categories. In year y+1, the area of a land use category is the sum of the area staying in the same land use category (e.g.,  $ca_{11y}$ ) plus the area that transitioned to that land use category from other land use categories (e.g.,  $ca_{21y}$ ,  $ca_{31y}$ , and  $ca_{i1y}$ ). Area of land use category *j* in year y+1,  $A_{jy+1}$ , is calculated as follows:

$$A_{j\,y+1} = \sum_{i} c a_{ij\,y}$$
 Equation 97

Where:

 $A_{j y+1}$ Area of land use category j in project area in year y+1; ha $ca_{ij y}$ Area of land converted from land use category i to j in the project area in year y;<br/>ha

Please refer to section J for determination of  $EF_{ij}$ ,  $A_{i0}$  and  $p_{ij}$ , i.e., MEF,  $MA_0$  and MP.

#### H. Calculation of project net emissions

Project net emissions in year *y* during the monitoring period are estimated based on results of monitoring, and calculated as follows:

$$PE_y = \Delta CS_{pjy} * 44/12 + E_{fuely} + E_{fertilizery} + DE_y$$

Equation 108

Where:

$PE_y$	Project net emissions in year y; tCO <sub>2</sub> -eq
$\Delta CS_{pjy}$	Carbon stock change in the project area in year y; tC
$E_{fuel \; y}$	$CO_2$ emissions from fossil fuel combustion in year y due to the project activities;
	tCO <sub>2</sub>
$E_{\text{fertilizer y}}$	GHG emissions from fertilizer application within the project area and the activity
	area as a part of the project activities in year y; tCO <sub>2</sub> -eq
$DE_y$	Displaced emissions to the displacement belt in year $y$ ; tCO <sub>2</sub>

(1) Carbon stock change in the project area in year y,  $\Delta CS_{pjy}$ 

The same option applied in the calculation of the project reference level must be used for the project emissions.

Option 1: Use the National FRL transition probabilities from forest to non-forest classes only

Carbon stock change in the project area in year <i>y</i> is calculated as follows:	
$\Delta CS_{pjy} = \sum_{i} \Delta CS_{pjiy} = \sum_{i} CA_{pjiy} * EF_{i}$	Equation <u>11</u> 9

Where:

$\Delta CS_{pj y}$	Carbon stock change in the project area in year y; tC
$\Delta CS_{pjiy}$	Carbon stock change in area converted from forest class $i$ to non-forest in the
	project area in year y; tC
CA <sub>pj i y</sub>	Area converted from forest class $i$ to non-forest in the project area in year $y$ ; ha
EFi	Emission factor applicable for forest class <i>i</i> ; tC ha <sup>-1</sup>

 $CA_{pjiy}$  will be determined using Cambodia's official forest maps provided by the government; calculated as the yearly average, if the monitoring interval is more than one year.

Accuracy assessment: to better account for the uncertainty that may be present in Cambodia's official forest map within the project area, an accuracy assessment will be performed of the project area and displacement belt. The accuracy assessment will be performed on the most recent official forest map and using a combination of the points used by the government of Cambodia and with additional points to ensure representative coverage of all classes within the project area. The results of the accuracy assessment of the project area and the displacement belt will determine the error-adjusted area estimates for each forest class. The accuracy assessment will be performed at the start of the project and each subsequent monitoring period to ensure high quality estimates of emission and reductions. This process is based on the methods included in Olofsson et al. 2014 and is endorsed by the FAO. However, the error

adjustment of the areas will not be performed, if the government of Cambodia decides to require the project to use Cambodia's official forest map as it is.

See section J for  $EF_i$ .

Option 2: Use all the National FRL transition probabilities among classes that would result in GHG emissions

To estimate the total emissions, the carbon stock change in the project area in year y is calculated as follows:

 $\Delta CS_{pjy} = \sum_{i} \sum_{j} cs_{pjijy} \begin{cases} cs_{pjijy}, & \text{if } cs_{pjijy} > 0\\ 0, & \text{otherwise} \end{cases}$  Equation <u>1240</u>

Where:

 $\Delta CS_{pjy}$  Carbon stock change in the project area in year y; tC

 $cs_{pj ij y}$  Carbon stock change in the project area from changes of land use category *i* to *j* in year *y*; tC

The notation  $cs_{pj \ ij \ y}$  is an element of *i*-by-*j* cross-tabulation matrix  $MCS_{pj \ y}$  which is a product of element-wise multiplication (Hadamard product) of *i*-by-*j* matrices of area changes and emission factors.

$$MCS_{pjy} = MCA_{pjy} \cdot MEF$$
 Equation 1311

Where:

$MCS_{pjy}$	i-by- $j$ matrix in which each element is carbon stock change in the project area
	from changes in land use category <i>i</i> to <i>j</i> in year <i>y</i> , $cs_{pj ij y}$ ; tC
MCA <sub>pj y</sub>	<i>i</i> -by- <i>j</i> matrix in which each element is the area of land converted from land use
	category <i>i</i> to <i>j</i> in the project area in year <i>y</i> , $ca_{pj \ ij \ y}$ ; ha
ca <sub>pj ij y</sub>	Area of land converted from land use category $i$ to $j$ in the project area in year $y$ ;
	ha
MEF	<i>i</i> -by- <i>j</i> matrix in which each element is emission factor for area of land converted
	from land use category <i>i</i> to <i>j</i> , $EF_{ij}$ ; tC ha <sup>-1</sup>

 $ca_{pj \ ij \ y}$  will be determined using Cambodia's official forest maps provided by the government; calculated as the yearly average, if the monitoring interval is more than one year.

To quantify and account for the uncertainty of Cambodia's official forest map in the project

area, an accuracy assessment will be performed as described above.

See section J for  $EF_{ij}$ .

#### (2) Emissions due to the project activities

Based on the requirements of the methodology guidelines ver01.0, emissions resulting from the implementation of the project activities shall be accounted, including fossil fuel combustion and the application of fertilizer (synthetic fertilizer, organic fertilizer, N-fixing crop, liming material and urea fertilizer). If the GHG emissions, due to use of fuel or the application of fertilizers, are estimated to less than five percent of the total net emissions at the time of validation, quantities of such GHG sources do not need to be monitored, and planned quantities can be used for calculation of GHG emissions.

#### Emissions from fossil fuel combustion

 $CO_2$  emissions from fossil fuel combustion resulting from the implementation of project activities<sup>1</sup> are calculated by applying the following direct method or indirect method, and these can be used interchangeably or simultaneously for different types of vehicle and equipment. E.g., project participant can calculate  $E_{fuel y}$  by applying the direct method for motorbikes and the in-direct method for trucks.

#### Direct method

$E_{fuel v} = \sum_{f} E_{fuel f v}$	$=\sum_{f} (FC_{fv})$	$*NCV_{f} *$	$EF_{fuel}$	$_{f})$
--------------------------------------	-----------------------	--------------	-------------	---------

Equation <u>1412</u>

Where:

$E_{\text{fuel y}}$	$CO_2$ emissions from fossil fuel combustion in year y due to the project activities;
	tCO <sub>2</sub>
$E_{fuel\;f\;y}$	$CO_2$ emissions from combustion of fossil fuel type <i>f</i> in year <i>y</i> ; t $CO_2$
$FC_{fy}$	Quantity of fuel type <i>f</i> consumed in year <i>y</i> ; kg
$NCV_{\mathrm{f}}$	Net calorific value of fuel <i>f</i> ; GJ kg <sup>-1</sup>
$\mathrm{EF}_{\mathrm{fuel}\;\mathrm{f}}$	$CO_2$ emission factor of the fuel type <i>f</i> combusted; t $CO_2$ GJ <sup>-1</sup>
f	fuel types combusted; dimensionless
Sec. costion	I for NCV and EE

See section J for *NCV<sub>f</sub>* and *EF*<sub>fuel f</sub>.

<sup>&</sup>lt;sup>1</sup> A/R methodological tool "Estimation of GHG emissions related to fossil fuel combustion in A/R CDM project activities" was referenced.

Indirect m	ethod	
$E_{fuel y} = 2$	$\sum_{f} \sum_{j} E_{fuel j f y} = \sum_{f} \sum_{j} (NVE_{j f y} * TDU_{j f y} * SEC_{j f} * NC$	$V_f * EF_{fuel f}$ )
		Equation <u>15</u> 13
Where:		
E <sub>fuel y</sub>	$CO_2$ emissions from fossil fuel combustion in year <i>y</i> due to the $tCO_2$	e project activities;
E <sub>fuel i f v</sub>	$CO_2$ emissions from fossil fuel combustion in vehicle or equ	ipment type <i>j</i> using
5.5	fuel type <i>f</i> in year <i>y</i> ; tCO <sub>2</sub>	
NVE <sub>jfy</sub>	Number of vehicle or equipment type $j$ using fuel type $f$ in ye	ar y; unit
TDU <sub>jfy</sub>	Total travel distance for vehicle type $j$ or use hours for equi	pment type <i>j</i> using
5.5	fuel type f in year y; km or hour unit <sup>-1</sup>	
SEC <sub>jf</sub>	Average specific energy consumption of vehicle or equipment	t type <i>j</i> for fuel type
5	f; kg km <sup>-1</sup> or hour <sup>-1</sup>	
$\mathrm{NCV}_{\mathrm{f}}$	Net calorific value of fuel $f$ ; GJ kg <sup>-1</sup>	
EF <sub>fuel f</sub>	$CO_2$ emission factor of the fuel type <i>f</i> combusted; t $CO_2$ GJ <sup>-1</sup>	
f	fuel types combusted; dimensionless	
j	type of vehicle or equipment; dimensionless	
For <i>NCV</i> <sub>f</sub> ,	$EF_{fuel f}$ , and $SEC_{jf}$ , see section J.	
Emissions	from fertilizer application	
GHG emis	sions from fertilizer application are calculated as follow:	
E <sub>fertilizer y</sub> =	Edirect-N y + Eindirect-N y + Eliming y + Eurea y	Equation <u>16</u> 14
Where:		
Efertilizer y	GHG emissions from fertilizer application within the project a	area and the activity
	area for implementation of the project activities in year y; tCC	D <sub>2</sub> -eq
Edirect-N y	Direct N <sub>2</sub> O emissions as a result of nitrogen application with	hin the project area
	and the activity area for implementation of the project activit	ies in year <i>y</i> ; tCO <sub>2</sub> -
	eq	
Eindirect-N y	Indirect N <sub>2</sub> O emissions as a result of nitrogen application wit	hin the project area
	and the activity area for implementation of the project activit	ies in year <i>y</i> ; tCO <sub>2</sub> -
	eq	
Eliming y	CO2 emissions as a result of adding liming materials within t	he project area and
	the activity area for implementation of the project activities in	n year $v$ ; tCO <sub>2</sub>

 $E_{urea y}$  CO<sub>2</sub> emissions as a result of urea fertilization application within the project area and the activity area for implementation of the project activities in year y; tCO<sub>2</sub>

Direct  $N_2O$  emissions as a result of nitrogen application for the implementation of the project activities are calculated as follow<sup>2</sup>:

 $E_{direct-Ny} = \sum_{c} [(F_{SN cy} + F_{ON cy} + F_{CR cy}) * EF_{direct-Nc}] * 44/28 * GWP_{N20}$ Equation <u>1715</u>

Where:

- $E_{direct-N y}$  Direct N<sub>2</sub>O emissions as a result of nitrogen application within the project area and the activity area for implementation of the project activities in year y; tCO<sub>2</sub>eq
- F<sub>SN c y</sub> Mass of nitrogen in synthetic fertilizer applied for implementation of the project activities in cropland type *c* in the project area and the activity area in year *y*; tN
   F<sub>ON c y</sub> Mass of nitrogen in organic fertilizer made from materials sourced from outside of the project area and the activity area and applied for implementation of the project activities in cropland type *c* in the project area and the activity area in year
- $F_{CR cy}$  Mass of nitrogen in crop residues (above-ground and below-ground) in N-fixing crops, introduced for implementation of the project activities in cropland type *c* in the project area and the activity area and returned to soils, in year *y*; tN
- $EF_{direct-N c}$  Emission factor for N<sub>2</sub>O emissions from nitrogen inputs in cropland type c; t N<sub>2</sub>O-N tN-input<sup>-1</sup>

44/28 Ratio of molecular weight of N<sub>2</sub>O and N; dimensionless

 $GWP_{N2O}$  Global Warming Potential for N<sub>2</sub>O; tCO<sub>2</sub> (t N<sub>2</sub>O)<sup>-1</sup>

c Types of croplands: upland cropland and flooded cropland such as rice paddy; dimensionless

See section J for  $EF_{direct-Nc}$  and  $GWP_{N2O}$ .

v; tN

 $F_{SN c y} = M_{SN c y} * NC_{SN c}$ Equation 1846 $F_{ON c y} = M_{ON c y} * NC_{ON c}$ Equation 1947Where: $F_{SN c y}$ Mass of nitrogen in synthetic fertilizer applied for implementation of the project

<sup>&</sup>lt;sup>2</sup> This is based on A/R Methodology tool "Estimation of direct nitrous oxide emission from nitrogen fertilization" and 2006 IPCC Guidelines for National Greenhouse Gas Inventories.

	activities in cropland type $c$ in the project area and the activity area in year $y$ , tN
Fon c y	Mass of nitrogen in organic fertilizer made from materials sourced from outside
	of the project area and the activity area and applied for implementation of the
	project activities in cropland type $c$ in the project area and the activity area in year
	y, tN
$M_{SNcy}$	Mass of synthetic fertilizer applied for implementation of the project activities in
	cropland type $c$ in the project area and the activity area in year $y$ , t
$M_{ONcy}$	Mass of organic fertilizer made from materials sourced from outside of the project
	area and the activity area and applied for implementation of the project activities
	in cropland type $c$ in the project area and the activity area in year $y$ , t
$NC_{SN c}$	Nitrogen content of synthetic fertilizer applied in cropland type <i>c</i> ; tN (t fertilizer)
	-1
NC <sub>ON c</sub>	Nitrogen content of organic fertilizer applied in cropland type $c$ ; tN (t fertilizer) <sup>-</sup>
	1

Data from producers of synthetic fertilizer and published data are used for  $NC_{SN c} NC_{ON c}$ , respectively. If multiple types of synthetic or organic fertilizers are used, choose the highest data as a conservative estimation. Note that organic fertilizer which is made from organic materials sourced from inside the project area and the activity area are NOT accounted because the emissions from those organic materials occur in the areas regardless the implementation of the project activities.

$$F_{CR\,c\,y} = \sum_{T} \left[ Crop_{c\,T\,y} * Area_{c\,T\,y} * Frac_{Renew\,T} * (R_{AG\,T} * N_{AG\,T} + R_{BG\,T} * N_{BG\,T}) \right]$$
Equation 2018

Where:

$F_{CR \ c \ y}$	Mass of nitrogen in crop residues (above-ground and below-ground) in N-fixing
	crops, introduced for implementation of the project activities in cropland type $c$
	in the project area and the activity area and returned to soils, in year y; tN
Crop <sub>c T y</sub>	Harvested annual dry matter yield for N-fixing crop $T$ per unit area, introduced
	for implementation of the project activities in cropland type $c$ in the project area
	and the activity area in year y; t d.m. ha $^{-1}$
Area <sub>cTy</sub>	Total annual area harvested of N-fixing crop $T$ , introduced for implementation of
	the project activities in cropland type $c$ in the project area and the activity area in
	year y; ha
$Frac_{Renew T}$	Fraction of total area under N-fixing crop $T$ that is renewed annually;
	dimensionless, 0-1

R <sub>AG T</sub>	Ratio of above-ground residues to harvested yield for N-fixing crop $T$ ; t d.m. (t
	d.m.) <sup>-1</sup>
N <sub>AG T</sub>	N content of above-ground residues for N-fixing crop $T$ ; t N (t d.m.) <sup>-1</sup>
$R_{BGT}$	Ratio of below-ground residues to harvested yield for N-fixing crop $T$ ; t d.m. (t
	d.m.) <sup>-1</sup>
$N_{BGT}$	N content of below-ground residues for N-fixing crop $T$ ; t N (t d.m.) <sup>-1</sup>
Т	Types of N-fixing crops; dimensionless

Where cropland is renewed on average every X years,  $Frac_{renew} = 1/X$ . For annual crops  $Frac_{renew} = 1$ .

Indirect  $N_2O$  emissions as a result of nitrogen application for implementation of the project activities are calculated as follow:

 $E_{indirect-N,y} = \left[ \left( F_{SN y} * Frac_{SN} + F_{ON y} * Frac_{ON} \right) * EF_{indirect-N} + \left( F_{SN y} + F_{ON y} + F_{CR y} \right) * Frac_{leach} * EF_{leach-N} \right] * \frac{44}{28} * GWP_{N20}$ Equation <u>2149</u>

Where:

$E_{indirect-Ny}$	Indirect N <sub>2</sub> O emissions as a result of nitrogen application within the project area
	and the activity area for implementation of the project activities in year $y$ ; tCO <sub>2</sub> -
	eq
$F_{SNy}$	Mass of nitrogen in synthetic fertilizer applied for implementation of the project
	activities in the project area and the activity area in year $y$ ; tN
$F_{ON \; y}$	Mass of nitrogen in organic fertilizer made from materials sourced from outside
	the project area and the activity area and applied for implementation of the project
	activities in the project area and the activity area in year $y$ ; tN
$F_{CR y}$	Mass of nitrogen in crop residues (above-ground and below-ground) in N-fixing
	crops, introduced for implementation of the project activities in the project area
	and the activity area and returned to soils, in year $y$ ; tN
Frac <sub>SN</sub>	Fraction that volatilized as $NH_3$ and $NO_X$ for synthetic fertilizers; dimensionless,
	0-1
Fracon	Fraction that volatilized as NH3 and NOX for organic fertilizers; dimensionless,
	0-1
EF <sub>indirect-N</sub>	Emission factor for $N_2O$ emissions from atmospheric deposition of N on soils and
	water surfaces; t N <sub>2</sub> O-N (t NH <sub>3</sub> -N and NO <sub>X</sub> -N volatilized) <sup>-1</sup>
Fracleach	Fraction of N that is lost through leaching and runoff; dimensionless, 0-1
$\mathrm{EF}_{\mathrm{leach-N}}$	Emission factor for $N_2O$ emissions from N leaching and runoff; t $N_2O\text{-}N$ (t N
	leaching and runoff) <sup>-1</sup>

44/28	Ratio of molecular weight of N2O and N; dimensionless
GWP <sub>N2O</sub>	Global Warming Potential for N <sub>2</sub> O; tCO <sub>2</sub> (t N <sub>2</sub> O) <sup>-1</sup>

See Section J for Frac<sub>SN</sub>, Frac<sub>ON</sub>, Frac<sub>leach</sub>, EF<sub>indirect-N</sub>, EF<sub>leach-N</sub> and GWP<sub>N2O</sub>.

$F_{SN y} = \sum_{c} F_{SN c y}$	Equation <u>22</u> 20
$F_{ONy} = \sum_{c} F_{ONcy}$	Equation 2321
$F_{CR y} = \sum_{c} F_{CR c y}$	Equation <u>24</u> 22

Where:

$F_{SNy}$	Mass of nitrogen in synthetic fertilizer applied for implementation of the project
	activities in the project area and the activity area in year $y$ ; tN
$F_{ON  y}$	Mass of nitrogen in organic fertilizer made from materials sourced from outside
	the project area and the activity area and applied for implementation of the project
	activities in the project area and the activity area in year y; tN
F <sub>CR y</sub>	Mass of nitrogen in crop residues (above-ground and below-ground) in N-fixing
	crops, introduced for implementation of the project activities in the project area
	and the activity area and returned to soils, in year $y$ ; tN
$F_{SNcy}$	Mass of nitrogen in synthetic fertilizer applied for implementation of the project
	activities in cropland type $c$ in the project area and the activity area in year $y$ ; tN
F <sub>ON cy</sub>	Mass of nitrogen in organic fertilizer made from materials sourced from outside
	the project area and the activity area and applied for implementation of the project
	activities in cropland type $c$ in the project area and the activity area in year $y$ ; tN
$F_{CRcy}$	Mass of nitrogen in crop residues (above-ground and below-ground) in N-fixing
	crops, introduced for implementation of the project activities in cropland type $c$
	in the project area and the activity area and returned to soils, in year $y$ ; tN

Use  $F_{SN\,c\,y},\,F_{ON\,c\,y}$  and  $F_{CR\,c\,y}$  calculated in Equations 16, 17 and 18.

CO<sub>2</sub> emissions as a result of adding liming materials for implementation of the project activities are calculated as follow:

 $E_{liming y} = (M_{limestone y} * EF_{limestone} + M_{dolomite y} * EF_{dolomite}) * \frac{44}{12}$  Equation 2523

Where:

 $E_{\text{liming y}}$  CO<sub>2</sub> emissions as a result of adding liming materials within the project area and the activity area during implementation of the project activities in year *y*; tCO<sub>2</sub>

$M_{\text{limestone y}}$	Mass of calcic limestone (CaCO <sub>3</sub> ) applied for implementation of the project
	activities in the project area and the activity area in year y; t
EFlimestone	Emission factor for limestone; t C (t limestone) <sup>-1</sup>
$M_{\text{dolomite y}}$	Mass of dolomite (CaMg(CO <sub>3</sub> ) <sub>2</sub> ) applied for implementation of the project
	activities in the project area and the activity area in year y; t
EF <sub>dolomite</sub>	Emission factor for dolomite; t C (t dolomite) <sup>-1</sup>
44/12	Ratio of molecular weight of CO <sub>2</sub> and C; dimensionless

See Section J for *EF*<sub>limestone</sub> and *EF*<sub>dolomite</sub>.

CO<sub>2</sub> emissions as a result of urea fertilization application for implementation of the project activities are calculated as follow:

 $E_{urea y} = M_{urea y} * EF_{urea} * \frac{44}{12}$  Equation <u>26</u>24

Where:

E <sub>urea y</sub>	$\mathrm{CO}_2$ emissions as a result of application of urea within the project area and the
	activity area for implementation of the project activities in year $y$ ; tCO <sub>2</sub>
$M_{urea \ y}$	Mass of urea fertilizer applied for implementation of the project activities in the
	project area and the activity area in year y; t
EF <sub>urea</sub>	Emission factor for urea; t C/t urea
44/12	Ratio of molecular weight of CO <sub>2</sub> and C; dimensionless

See Section J for *EF*<sub>urea</sub>.

(3) Displaced emissions

In this methodology, displacement is assessed through monitoring of the displacement belt. The displacement belt captures the displacement of baseline activities due to the project activities, or interventions, in the project area.

Although a detailed analysis of the local drivers of deforestation and degradation is not anticipated, the local context will be captured through inputs from local and national experts including authorities. Displaced emissions are calculated as increases of emissions compared to reference emissions from the displacement belt which is separately calculated from reference emissions for the project reference level.

Displaced emissions are calculated as follows:

$DE_{\mathcal{Y}} = \begin{cases} DP_{\mathcal{Y}} \\ \end{cases}$	$A_{y} - DR_{y}$ , if $DP_{y} - DR_{y} > 0$ 0, otherwise	Equation <u>27</u> 25
Where:		
DEy	Displaced emissions to the displacement belt in year $y$ ; tCO <sub>2</sub>	
DR <sub>y</sub>	Reference emissions from the displacement belt in year $y$ ; tCO <sub>2</sub>	
DPy	Project emissions from the displacement belt in year $y$ ; tCO <sub>2</sub>	
$DR_y$ and $DP_y$	y are calculated as follows:	
$DR_y = \Delta CS_d$	<sub>y</sub> * 44/12	Equation <u>28</u> 26
Where:		
DR <sub>y</sub>	Reference emissions from the displacement belt in year $y$ ; tCO <sub>2</sub>	
$\Delta CS_{dy}$	Projected carbon stock change in the displacement belt in year y;	tC
44/12	Conversion factor of molecular weight of carbon to CO <sub>2</sub> ; dimension	sionless
$DP_y = \Delta CS_d$	<sub>pjy</sub> * 44/12	Equation <u>29<del>27</del></u>
Where:		
DPy	Project emissions from the displacement belt in year $y$ ; tCO <sub>2</sub>	
$\Delta CS_{dpjy}$	Actual carbon stock change in the displacement belt in year y; to	
44/12	Conversion factor of molecular weight of carbon to CO <sub>2</sub> ; dimension	sionless
Projected and actual carbon stock changes in the displacement belt area, i.e., $\Delta CS_{dy}$ and $\Delta CS_d$ <sub>pj y</sub> , are calculated by applying the option used in Section G.2 Calculation of project reference level.		
Option 1: U	se the National FRL transition probabilities from forest to non-fore	est classes only
Projected ca For the proje	rbon stock change in the displacement belt in year y is calculated a ect start year:	as follows:
$\Delta CS_{dy} = \sum_{i} A_{di0} * P_{di} * d_{y} / d_{0y} * EF_{i}$ Equation 3028		
For subsequent years to the project start year		
$\Delta CS_{dy} = \sum_{i} A_{diy-1} * P_{di} * \frac{d_y}{d_0y} * EF_i$ Equation <u>31</u> 28		
Where: $\Delta CS_{dy}$	Projected carbon stock change in the displacement belt in year v	tC

<u>Adio</u>	Area of forest class <i>i</i> in the displacement belt at the inception of the project; ha
$A_{diy\underline{-1}}$	Area of forest class $i$ in the displacement belt in year $y$ - $l$ ; ha
$P_{di}$	Annual transition probability from forest class $i$ to non-forest within the
	displacement belt; dimensionless, 0-1
<u>d</u> y	Number of operating days in year y, days
<u>d</u> o y	Number of days in year y, days
EFi	Emission factor applicable for forest class $i$ ; tC ha <sup>-1</sup>
Area of fore	st class $i$ is assumed to decrease every year due to deforestation, and therefore
calculated as	follows:
For the proje	tet start year:
$\underline{A}_{diy} = \underline{A}_{di0}$	* $(1 - P_{di}) * a_y / a_{0y}$ Equation 3229
For subseque	ent years to the project start year:
$A_{d i y+1} = A_{d i}$	$y_{-1} * (1 - P_{d_i})$ Equation 3329
Where:	
Adiv	Area of forest class <i>i</i> in the displacement belt in year <i>v</i> ; ha
Adio	Area of forest class <i>i</i> in the displacement belt at the inception of the project; ha
Adiv-1	Area of forest class <i>i</i> in the displacement belt in year $v$ - $l$ ; ha
P <sub>di</sub>	Annual transition probability from forest class <i>i</i> to non-forest within the
	displacement belt; dimensionless, 0-1
d <sub>v</sub>	Number of operating days in year y, days
$\underline{d}_{ov}$	Number of days in year y, days
Please refer	section J for $A_{di0}$ and $P_{di}$ .
Actual carbo	In stock change in the displacement belt in year $y$ is calculated as follows:
$\Delta CS_{d pj y} = \sum$	CA <sub>d pj i y</sub> * EF <sub>i</sub> Equation <u>34</u> 30
Where:	
$\Delta CS_{d  pj  y}$	Actual carbon stock change in the displacement belt in year y; tC
CA <sub>d pj i y</sub>	Area converted from forest class <i>i</i> to non-forest in the displacement belt in year
	y; ha
EFi	Emission factor applicable for land use category $i$ ; tC ha <sup>-1</sup>
$CA_{dpity}$ will be determined using Cambodia's official forest maps provided by the government:	
calculated as yearly average, if the monitoring interval is more than one year. Where evidence	

can be collected that deforestation in the displacement belt is not attributable to the project, the detected deforestation is not considered as displacement and therefore is excluded from  $CA_{d pj}$ *iy.* 

To quantify and account for the uncertainty of Cambodia's official forest map in the project area, an accuracy assessment will be performed as described in Section H calculation of project net emission, sub-section (1) carbon stock change in the project area.

See section J for  $EF_i$ .

Option 2: Use all transition probabilities resulting in emissions in the National FRL

<u>Projected carbon stock change</u> in the displacement belt in year y is calculated as follows: To estimate the total emissions, the projected carbon stock change in year y in the displacement belt is calculated as follows:

$$\Delta CS_{dy} = \sum_{i} \sum_{j} cs_{dijy} \begin{cases} cs_{dijy}, & \text{if } cs_{dijy} > 0\\ 0, & \text{otherwise} \end{cases}$$
 Equation 3531

Where:

 $\Delta CS_{dy}$ Projected carbon stock change in the displacement belt in year y; tC $cs_{dijy}$ Projected carbon stock change in the displacement belt from changes of land use<br/>category i to j in year y; tC

The notation  $cs_{dij}$  is an element of *i*-by-*j* cross-tabulation matrix  $MCS_{dy}$  which is a product of element-wise multiplication (Hadamard product) of *i*-by-*j* matrices of area changes and emission factors.

$$MCS_{dv} = MCA_{dv} \cdot MEF$$

Equation <u>36</u>32

Where:

$MCS_{dy}$	<i>i</i> -by- <i>j</i> matrix in which each element is projected carbon stock change in the
	displacement belt from changes of land use category <i>i</i> to <i>j</i> in year <i>y</i> , $cs_{dijy}$ ; tC

- MCA<sub>d y</sub> i-by-j matrix in which each element is projected area of land converted from land use category i to j in the displacement belt in year y,  $ca_{d ij y}$ ; ha
- MEF *i*-by-*j* matrix in which each element is the emission factor for area of land converted from land use category *i* to *j*,  $EF_{ij}$ ; tC ha<sup>-1</sup>

Projected area of land conversion in year *y* is calculated as the product between areas of land categories in the displacement belt in year *y* and annual transition probability.

Equation 4036

 $MCA_{dy} = MA_{dy-1}MP_d$ Equation 3733 Where: MCA<sub>d v</sub> *i*-by-*j* matrix in which each element is the projected area of land converted from land use category *i* to *j* in the displacement belt in year *y*,  $ca_{dij}$ , ha *i*-by-*j* diagonal matrix whose diagonal elements are areas of land use categories MAd y-1 in the displacement belt in year y-1,  $A_{diy-l}$ ; ha MP<sub>d</sub> *i*-by-*j* matrix in which each element is the annual transition probability from land use category *i* to *j* in the displacement belt,  $p_{dij}$ ; dimensionless, 0-1 Similar to Equation 7, area of land use category j in year y+1,  $A_{djy+1}$ , is calculated as follows:  $A_{d\,j\,y+1} = \sum_i c a_{d\,ij\,y}$ Equation 3834 Where: Area of land use category *j* in the displacement belt in year y+1; ha  $A_{d j y+1}$ Area of land converted from land use category *i* to *j* in the displacement belt in ca<sub>d ij y</sub> year y; ha Please refer section J for determination of  $EF_{ij}$ ,  $A_{di0}$  and  $p_{dij}$ , i.e., MEF,  $MA_{d0}$  and  $MP_d$ . Actual carbon stock change in the displacement belt in year y is calculated as follows: To estimate the total emissions, the carbon stock change in the displacement belt in year y is calculated as follows:  $\Delta CS_{d p j y} = \sum_{i} \sum_{j} cs_{d p j i j y} \begin{cases} cs_{d p j i j y}, & \text{if } cs_{d p j i j y} > 0\\ 0, & \text{otherwise} \end{cases}$ Equation 3935

Where:

 $\Delta CS_{d pj y} \qquad \text{Actual carbon stock change in the displacement belt in year } y; tC$   $cs_{d pj ij y} \qquad \text{Carbon stock change in the displacement belt from changes of land use category}$  i to j in year y; tC

The notation  $cs_{d pj ij}$  is an element of the *i*-by-*j* cross-tabulation matrix  $MCS_{d pj y}$  which is a product of element-wise multiplication (Hadamard product) of *i*-by-*j* matrices of area changes and emission factors.

$$MCS_{d p j y} = MCA_{d p j y} \cdot MEF$$

Where:

$MCS_{dpjy}$	<i>i</i> -by- <i>j</i> matrix in which each element is carbon stock change in the displacement
	belt from changes of land use category <i>i</i> to <i>j</i> in year <i>y</i> , $cs_{d pj ij y}$ ; tC
$MCA_{d\ pj\ y}$	<i>i</i> -by- <i>j</i> matrix in which each element is area of land converted from land use
	category <i>i</i> to <i>j</i> in the displacement belt in year <i>y</i> , $ca_{d p j i j y}$ ; ha
ca <sub>d pj ij y</sub>	Area of land converted from land use category $i$ to $j$ in the displacement belt in
	year y; ha
MEF	<i>i</i> -by- <i>j</i> matrix in which each element is emission factor for area of land converted
	from land use category <i>i</i> to <i>j</i> , $EF_{ij}$ ; tC ha <sup>-1</sup>

 $ca_{d\ pj\ ij\ y}$  will be determined by using Cambodia's official forest maps provided by the government, will be calculated as yearly average, if the monitoring interval is more than one year. Where evidence can be collected that deforestation in the displacement belt is not attributable to the project, the detected deforestation is not considered as displacement and therefore is excluded from  $MCA_{d\ pj\ y}$ .

To quantify and account for the uncertainty of Cambodia's official forest map in the project area, an accuracy assessment will be performed as described in Section H calculation of project net emission, sub-section (1) carbon stock change in the project area.

See section J for  $EF_{ij}$ .

### I. Calculation of project emission reductions to be credited

Project emission reductions in year y are calculated as the difference between the project reference level and the project net emissions. Equation <u>41</u>37  $ER_v = RL_v - PE_v$ Where:  $ER_v$ Project emissions reductions in year y; tCO<sub>2</sub>-eq  $RL_y$ Project reference level in year y; tCO<sub>2</sub>  $PE_v$ Project net emissions in year y; tCO<sub>2</sub>-eq  $ER_{credt y} = ER_y * (1-DF)$ Equation 4238 Where: Project emissions reductions available to be credited in year y; tCO<sub>2</sub>-eq ER<sub>credit y</sub>,

ERyProject emissions reductions in year y; tCO2-eqDFDiscount factor, default as 0.2

A discount factor is applied as a measure for risk of reversals.

Project emissions reductions available to be credited for a monitoring period are calculated as follows:

 $ER_{credit p} = \sum_{i} ER_{credit y}$ 

Equation <u>43</u>39

Where:

 ER<sub>credit p</sub> Project emissions reductions available to be credited for a monitoring period *p*; tCO<sub>2</sub>-eq
 ER<sub>credit y</sub>, Project emissions reductions available to be credited in year *y*; tCO<sub>2</sub>-eq

### J. Data and parameters fixed ex ante

The lists of the source of each data and parameter fixed *ex ante* and the source of each of default values selected in this methodology are available at the end of this section. The same option applied in the calculation of project reference level must be used.

(1) Emission factors for land conversions

Emission factors are calculated by applying carbon stock data used in National FRL, or other official report by Cambodian government.

<u>Table 1</u> Table 1 shows carbon stock data used in the National FRL submitted in 2017. As new official data becomes available, emission factors can be updated in coordination with government of Cambodia.

Table 1 Above ground and below ground biomass data used in the National FRL submitted in

2017							
Forest/Non-	Land use category	Abbreviation	Above-	Below-	Total	biomass,	
Forest			ground,	ground,	tCha <sup>-1</sup>		
			tCha <sup>-1</sup>	tCha <sup>-1</sup>			
Forest	Evergreen forest	Е	76.61	14.69		91.30	
	Semi-evergreen	SE	114.21	20.9		135.11	
	forest						

	Pine forest	Р	47	9.54	56.54
	Deciduous forest	D	39.95	8.26	48.21
	Bamboo	В	0	0	0
	Mangrove	М	70.5	13.65	84.15
	Rear Mangrove	MR	77.55	14.85	92.40
	Flooded forest	FF	32.9	6.96	39.86
	Forest regrowth	FR	35.25	7.4	42.65
	Tree plantation	ТР	47	9.54	56.54
	Pine plantation	РР	47	9.54	56.54
Non-forest	Non-forest	NF	0	0	0

Although Cambodia defined tree plantation (TP) and pine plantation (PP) as forest, emissions and removals in areas converted from the other forest categories to these two categories were excluded from the National FRL submitted in 2017 in consideration of safeguards, while those in areas converted from TP and PP to the other land use categories are included. The project participant uses the latest National FRL, and apply zero as EF for the conversion categories which are excluded from calculation in the National FRL.

Option 1: Use the National FRL transition probabilities from forest to non-forest classes only  $EF_i$  is an emission factor for area of land converted from land use category *i*, which is forest, to non-forest.

$$EF_i = \sum_k C_{ik}$$

Equation 4440

Where:

EFi	Emission factor applicable for forest class $i$ ; tC ha <sup>-1</sup>
C <sub>ik</sub>	Carbon stock in carbon pool k in forest class i per unit area; tC ha <sup>-1</sup>
k	Carbon pools included in establishment of National FRL; dimensionless

Above ground and below ground are the carbon pools included in the establishment of the National FRL submitted in 2017. <u>Table 2</u> shows the calculated  $EF_i$  based on the National FRL submitted in 2017.

Table 2 Emission factors, EF<sub>i</sub>, for Option 1 based on the National FRL submitted in 2017

Forest class, <i>i</i>	Abbreviation	EF <sub>i</sub> , tC ha <sup>-1</sup>
Evergreen forest	Е	91.30

Semi-evergreen forest	SE	135.11
Pine forest	Р	56.54
Deciduous forest	D	48.21
Bamboo	В	0
Mangrove	М	84.15
Rear Mangrove	MR	92.40
Flooded forest	FF	39.86
Forest regrowth	FR	42.65
Tree plantation	ТР	56.54
Pine plantation	PP	56.54
Non-forest	NF	0

Option 2: Use all the National FRL transition probabilities among classes that would result in emissions

 $EF_{ij}$  is an emission factor for area of land converted from land use category *i* to *j*.

$$EF_{ij} = \sum_k C_{ik} - \sum_k C_{jk}$$

Equation 4541

Where:

EF <sub>ij</sub>	Emission factor for area of land converted from land use category $i$ to $j$ ; tC ha <sup>-1</sup>
$C_{ik}$	Carbon stock in carbon pool k in land use category i per unit area; tC ha <sup>-1</sup>
k	Carbon pools included in establishment of National FRL; dimensionless

Table 3 Emission factors,  $EF_{ij}$ , for Option 2 based on the National FRL submitted in 2017

		Land use category j											
		Е	SE	Р	D	В	М	MR	FF	FR	TP	PP	NF
i	Е	0.00	NA	34.76	43.09	91.30	7.15	NA	51.44	48.65	NA	NA	91.30
	SE	43.81	0.00	78.57	86.90	135.11	50.96	42.71	95.25	92.46	NA	NA	135.11
	Р	NA	NA	0.00	8.33	56.54	NA	NA	16.68	13.89	NA	NA	56.54
	D	NA	NA	NA	0.00	48.21	NA	NA	8.35	5.56	NA	NA	48.21
	В	NA	NA	NA	NA	0.00	NA	NA	NA	NA	NA	NA	0.00
	М	NA	NA	27.61	35.94	84.15	0.00	NA	44.29	41.50	NA	NA	84.15
	MR	1.10	NA	35.86	44.19	92.40	8.25	0.00	52.54	49.75	NA	NA	92.40
	FF	NA	NA	NA	NA	39.86	NA	NA	0.00	NA	NA	NA	39.86
	FR	NA	NA	NA	NA	42.65	NA	NA	2.79	0.00	NA	NA	42.65
	TP	NA	NA	0.00	8.33	56.54	NA	NA	16.68	13.89	0.00	0.00	56.54
	PP	NA	NA	0.00	8.33	56.54	NA	NA	16.68	13.89	0.00	0.00	56.54
	NF	NA	NA	NA	NA	0.00	NA	NA	NA	NA	NA	NA	0.00

Note that when a land use category with a higher carbon stock is converted to a lower carbon stock, e.g., conversion from evergreen forest to non-forest, the  $EF_{ij}$  is positive. Removals are not included in this methodology, and therefore emission factors for conversions from a land

use category with lower carbon stock to higher carbon stock are not available in <u>Table 3</u>Table 3. As previously described, emissions and removals in areas converted from the other forest categories to tree plantation (TP) or pine plantation (PP) were excluded in the establishment of the National FRL submitted in 2017, and therefore *EFs* of those transitions as well as transitions resulting in removals are also not available in <u>Table 3</u>Table 3.

(2) Area of land use category *i* at the inception of the projectThis method applies to both options, Option 1 and 2, and to the following parameters:

- $A_{i0}$  Area of forest class *i* or land use category *i* in the project area at the inception of the project; ha
- $A_{d\,i\,0}$  Area of forest class *i* or land use category *i* in the displacement belt at the inception of the project; ha

The most recent version of Cambodia's official forest map ( $\leq 2$  years) provided by Cambodian government used to determine the area of land use category *i* in the project area and in the displacement belt at the project start date. Once  $A_{i0}$  and  $A_{di0}$  are determined in the project design document, these areas, determined at the project start date, will not be updated.

To quantify and account for the uncertainty of Cambodia's official forest map in the project area, an accuracy assessment will be performed as described in Section H calculation of project net emission, sub-section (1) carbon stock change in the project area.

(3) Annual transition probability in the reference area

Option 1: Use the National FRL transition probabilities from forest to non-forest classes only Annual deforestation rate for each forest type existing inside the project area,  $P_i$ , is calculated by applying area,  $A_{refi}$ , and area change,  $CA_{refi}$ , from the National FRL and nationally endorsed data sets, and is used in Equation 2. <u>Table 4</u> Table 4 shows  $P_i$  calculated based on the National FRL submitted in 2017.

$$P_i = CA_{ref i} / (A_{ref i} * T_{ref})$$

Equation 4642

Where:

$\mathbf{P}_{i}$	Annual transition probability from forest class <i>i</i> to non-forest within the reference
	area; dimensionless, 0-1
CA <sub>ref i</sub>	Area converted into non-forest from forest class $i$ during the reference period,
	e.g., 2006-2014 in the National FRL submitted in 2017, in reference area; ha
A <sub>ref i</sub>	Area of forest class $i$ in the reference area at the first year, e.g., 2006 in the

National FRL submitted in 2017, of the reference period; ha

 $T_{ref}$ 

Number of years, e.g., eight in the National FRL submitted in 2017, in the reference period; year

Forest class	Abbreviation	P <sub>i</sub> , dimensionless
Evergreen forest	Е	0.0249
Semi-evergreen forest	SE	0.0309
Pine forest	Р	0.0000
Deciduous forest	D	0.0345
Bamboo	В	0.0141
Mangrove	М	0.0100
Rear Mangrove	MR	0.0417
Flooded forest	FF	0.0506
Forest regrowth	FR	0.0972
Tree plantation	ТР	0.1169
Pine plantation	РР	0.000

Table 4 P<sub>i</sub> calculated based on the data used in the National FRL submitted in 2017

Option 2: Use all the National FRL transition probabilities among classes that would result in GHG emissions

Annual transition probability from land use category *i* to *j* is obtained in the form of a transition probability matrix, MP, using the forest area change matrix reported in National FRL and nationally endorsed data sets. It is important to note that the annual transition probability cannot be obtained by simply dividing transition probability in multiple years by number of the years, because transition probability in multiple years, *c* years, is *c*-th power of annual transition probability. It is recommended to use a method provided in a peer reviewed scientific paper for calculation of the annual transition probability, such as Takada et al. 2010<sup>3</sup>. Calculated MP is used in Equation 6.

Obtain an area-based *i*-by-*j* transition matrix for each of time interval from the National FRL. In the National FRL submitted in 2017, Forest Area Change Matrices between 2006 and 2010 and between 2010 and 2014 area are available in Annex I Emission/Removal Calculation Tables. Annual transition probabilities derived using the matrices in the National FRL submitted in 2017 are available in Table 6Table 6 below.

<sup>&</sup>lt;sup>3</sup> Takada T, Miyamoto A and Hasegawa SF (2010) Derivation of a yearly transition probability matrix for land-use dynamics and its applications. Landscape Ecol 25:561-572

Transition probability in an observation interval t is calculated as follows: $prob_{ij t} = CA_{ref ij t}/A_{ref i t}$ Equation 4743

Where:

prob <sub>ij t</sub>	Transition probability from land use category $i$ to $j$ during an observation time
	interval <i>t</i> ; dimensionless, 0-1
CA <sub>ref ij t</sub>	Area converted from land use category $i$ to $j$ during an observation time interval
	t within the reference area; ha
A <sub>refit</sub>	Area of land use category $i$ at the first year of an observation time interval $t$ within
	reference area; ha

The notion  $prob_{ij t}$  is an element of *i*-by-*j* cross-tabulation matrix  $MP_{multi t}$ , and  $MP_{multi t}$  is the  $T_{multi}$ -th power of annual transition matrix  $MP_t$ .

$$MP_{multit} = MP_t^{T_{multi}}$$

Equation <u>48</u>44

Where:

 $\begin{array}{ll} \text{MP}_{\text{multi t}} & i\text{-by-}j \text{ matrix in which each element is the transition probability from land use} \\ & \text{category } i \text{ to } j \text{ during an observation time interval } t, prob_{ij t}; \text{ dimensionless, 0-1} \\ \text{MP}_{\text{t}} & i\text{-by-}j \text{ matrix in which each element is the annual transition probability from land} \\ & \text{use category } i \text{ to } j \text{ during an observation time interval } t, p_{ij t}; \text{ dimensionless, 0-1} \\ & \text{T}_{\text{multi}} & \text{number of years of observation interval; year} \end{array}$ 

<u>Table 5</u> shows the transition probability matrix,  $MP_{multi \ 2006-2010}$ , in which each element  $prob_{ij \ 2006-2010}$  is the transition probability from *i* to *j* in 4 years between 2006 and 2010 in the cross-tabulation based on the National FRL submitted in 2017. Note that the diagonal elements represent the probabilities of land being in the same category.

Table 5 Cross-tabulation matrix of 4-year, 2006-2010	, transition probability based on the National
FRL submitted	in 2017

	2010												
2006		Е	SE	Р	D	В	М	MR	FF	FR	TP	PP	NF
	E	0.960	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.001	0.000	0.000	0.039
	Se	0.000	0.952	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000	0.000	0.045
	Р	0.000	0.000	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	D	0.000	0.000	0.000	0.971	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.029
	В	0.000	0.000	0.000	0.000	0.992	0.000	0.000	0.000	0.000	0.000	0.000	0.008
	М	0.004	0.002	0.000	0.000	0.000	0.968	0.001	0.000	0.000	0.000	0.000	0.025
	MR	0.000	0.003	0.000	0.000	0.000	0.003	0.962	0.000	0.002	0.000	0.000	0.030
	FF	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.790	0.001	0.000	0.000	0.209

FR	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.881	0.001	0.000	0.116
TP	0.000	0.000	0.000	0.001	0.000	0.000	0.000	0.000	0.004	0.184	0.000	0.811
PP												
NF	0.002	0.001	0.000	0.003	0.000	0.000	0.000	0.007	0.007	0.001	0.000	0.979

Obtain the annual transition probability matrix,  $MP_t$ , in which each element  $p_{ijt}$  is the annual transition probability during an observation time interval *t*, by using  $MP_{multit}$  and a method provided in a peer reviewed scientific paper for calculation of annual transition probability such as Takada et al. 2010. The obtained  $MP_t$  should be validated by applying Equation 44.

If multiple  $MP_t$ s from more than one observation time interval are derived, calculate average annual transition probability as follows:

$$p_{ij} = \frac{\sum_{t} p_{ij\,t}}{N_{int}}$$
Equation 4945

Where:

$p_{ij}$	Annual transition probability from land use category $i$ to $j$ within the reference
	area; dimensionless, 0-1

 $p_{ijt}$  Annual transition probability from land use category *i* to *j* during an observation time interval *t*; dimensionless, 0-1

N<sub>int</sub> Number of observation intervals reported in National FRL; interval

<u>Table 6</u> shows the annual transition probability matrix, MP, in which each element  $p_{ij}$  is annual transition probability from *i* to *j* based on observations during intervals of 2006-2010 and 2010-2014 presented in the National FRL submitted in 2017.

Table 6 Annual transition probability matrix based on observations during intervals of 2006-2010 and 2010-2014 presented in the National FRL submitted in 2017

	y+1												
у		Е	SE	Р	D	В	М	MR	FF	FR	ТР	PP	NF
	Е	0.971	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000	0.000	0.027
	SE	0.000	0.963	0.000	0.000	0.000	0.000	0.000	0.000	0.002	0.000	0.000	0.035
	Р	0.000	0.000	1.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	D	0.000	0.000	0.000	0.960	0.000	0.000	0.000	0.000	0.000	0.001	0.000	0.040
	В	0.000	0.000	0.000	0.000	0.985	0.000	0.000	0.000	0.001	0.000	0.000	0.015
	М	0.001	0.000	0.000	0.000	0.000	0.988	0.001	0.000	0.000	0.000	0.000	0.010
	MR	0.000	0.000	0.000	0.000	0.000	0.003	0.944	0.000	0.005	0.001	0.000	0.048
	FF	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.940	0.001	0.000	0.000	0.060
	FR	0.004	0.002	0.000	0.000	0.000	0.000	0.000	0.000	0.875	0.002	0.000	0.117
	TP	0.000	0.000	0.000	-0.001	0.000	0.000	0.000	-0.001	0.003	0.780	0.000	0.219
	PP	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	1.000	0.000
	NF	0.001	0.001	0.000	0.003	0.000	0.000	0.000	0.002	0.002	0.000	0.000	0.990

(4) Annual transition probability in the displacement belt

The same method as (3), Annual transition probability in the reference area, is applied for the

displacement belt instead of the reference area.

(5) Parameters for calculating emissions due to the project activities <u>Fossil fuel combustion</u>

Default net caloric value and net calorific based emission factors are available in the 2006 IPCC Guidelines, and those of selected fuels types are listed in the table for "the source of each data and parameter fixed *ex ante*".

Reference figures such as manufacturer specifications can be used for  $SEC_{jf}$ . If no specific energy consumption data are available, fuel consumption and distance and/or hours are recorded before the initial verification to calculate  $SEC_{jf}$ . Those figures can be used for the entire project period.

### Fertilizer application

Data and parameter fixed *ex ante* and default values for calculating  $N_2O$  and  $CO_2$  emissions are shown in the tables for "the source of each data and parameter fixed *ex ante*" and "the source of each of default values selected in this methodology" below.

Parameter	Description of data	Source
Aio	Area of forest class $i$ or land use category $i$ in the	Cambodia's official
	project area at the inception of the project; ha	forest map
Pi	Option (1) Annual transition probability from forest	Cambodia's official
	class <i>i</i> to non-forest within the reference area;	forest reference level
	dimensionless, 0-1	(National FRL)
<u>d</u> <sub>y</sub>	Number of operating days in year y	Decided based on
		starting date of project
		operation and expected
		operational lifetime of
		project
<u>d</u> oy	Number of days in year y	Automatically decided
		by number of year y
p <sub>ij</sub>	Option (2) Annual transition probability from land	Cambodia's official
	use category $i$ to $j$ within the reference area;	forest reference level
	dimensionless, 0-1	(National FRL)
$\overline{EF_i}$	Option (1) Emission factor applicable for forest	Cambodia's official
	class $i$ ; tC ha <sup>-1</sup>	forest reference level

The source of each data and parameter fixed *ex ante* is listed as below.

		(National FRL)					
EF <sub>ij</sub>	Option (2) Emission factor for area of land	Cambodia's official					
	converted from land use category <i>i</i> to <i>j</i> ; tC ha <sup>-1</sup>	forest reference level					
		(National FRL)					
$A_{di0}$	Area of forest class <i>i</i> in the displacement belt at the Cambodia's official						
	inception of the project; ha forest map						
P <sub>d i</sub>	Option (1) Annual transition probability from forest	Cambodia's official					
	class $i$ to non-forest within the displacement belt;	forest maps					
	dimensionless, 0-1						
p <sub>d ij</sub>	Option (2) Annual transition probability from land	Cambodia's official					
	use category $i$ to $j$ within the displacement belt;	forest maps					
	dimensionless, 0-1						
	Net calorific value of fuel $f$ ; GJ kg <sup>-1</sup>	Table 1.2 of Ch. 1 Vol. 2					
NCV	- Gas/Diesel oil: 43.0 TJ Gg <sup>-1</sup>	of 2006 IPCC					
	– Motor Gasoline: 44.3 TJ Gg <sup>-1</sup>	Guidelines					
	- Crude Oil: 42.3 TJ Gg <sup>-1</sup>	Guidennes					
	$CO_2$ emission factor of the fuel type <i>f</i> combusted; t	Table 3.2.1 of Ch. 3 and					
	$CO_2 GJ^{-1}$	Table 2.5 of Ch 2 Vol 2					
EF <sub>fuel f</sub>	- Gas/Diesel Oil: 74,100 kg $CO_2$ TJ <sup>-1</sup>	of 2006 IPCC Guidelines					
	- Motor Gasoline: $69,300 \text{ kg CO}_2 \text{ TJ}^{-1}$	Tables 2.5 and 3.2.1					
	<ul> <li>Crude Oil: 73,300 kg TJ<sup>-1</sup></li> </ul>	Tables 2.3 and 3.2.1					

The source of each of default values selected in this methodology is listed as below.

Parameter	Description of data	Source
	Emission factor for N <sub>2</sub> O emissions from N inputs in	
	cropland type $c$ ; tN <sub>2</sub> O-N (tN-input) <sup>-1</sup>	Table 11.1 of Ch. 11
$EF_{direct-N c}$	- Cropland in general: 0.01 tN <sub>2</sub> O-N (tN-input) <sup>-1</sup>	Vol. 4 of 2006 IPCC
	- Rice paddy (flooded rice field): 0.003 tN <sub>2</sub> O-N	Guidelines
	(tN-input) <sup>-1</sup>	
	Emission factor for N <sub>2</sub> O emissions from atmospheric	
	deposition of N on soils and water surfaces; $tN_2O$ -N (t	Table 11.3 of Ch. 11
$EF_{indirect-N}$	NH <sub>3</sub> -N and NO <sub>X</sub> -N volatilized) <sup>-1</sup>	Vol. 4 of 2006 IPCC
	- 0.010 t N <sub>2</sub> O-N (t NH <sub>3</sub> -N and NO <sub>X</sub> -N volatilized) <sup>-</sup>	Guidelines
	1	
EF <sub>leach-N</sub>	Emission factor for N <sub>2</sub> O emissions from N leaching	Table 11.3 of Ch. 11
	and runoff; $tN_2O$ -N (t leaching and runoff) <sup>-1</sup>	Vol. 4 of 2006 IPCC

	- 0.0075 tN <sub>2</sub> O-N (t N leaching and runoff) <sup>-1</sup>	Guidelines
EFlimestone	Emission factor for limestone; tC (t limestone) <sup>-1</sup>	Section 11.3.1 of Ch.
	- 0.12 tC (t limestone) <sup>-1</sup>	11 Vol. 4 of 2006
		IPCC Guidelines
$EF_{dolomite}$	Emission factor for dolomite; tC (t dolomite) <sup>-1</sup>	Section 11.3.1 of Ch.
	- 0.13 tC (t dolomite) <sup>-1</sup>	11 Vol. 4 of 2006
		IPCC Guidelines
$\mathrm{EF}_{\mathrm{urea}}$	Emission factor for urea; tC (t urea) <sup>-1</sup>	Section 11.4.1 of Ch.
	- 0.20 tC (t urea) <sup>-1</sup>	11 Vol. 4 of 2006
		IPCC Guidelines
GWP <sub>N20</sub>	Global Warming Potential for N <sub>2</sub> O; tCO <sub>2</sub> (tN <sub>2</sub> O) <sup>-1</sup>	Table 2.14 in Ch.2 of
	$-  GWP_{100} \text{ for } N_2O: 298 \text{ tCO}_2 \text{ (tN}_2O)^{-1}$	Working Group I
		contribution to the
		IPCC Forth
		Assessment Report
Frac <sub>sn</sub>	Fraction that volatilized as $NH_3$ and $NO_X$ for synthetic	Table 11.3 of Ch. 11
	fertilizers; dimensionless, 0-1	Vol. 4 of 2006 IPCC
	- 0.10	Guidelines
Fracon	Fraction that volatilized as NH <sub>3</sub> and NO <sub>X</sub> for organic	Table 11.3 of Ch. 11
	fertilizers; dimensionless, 0-1	Vol. 4 of 2006 IPCC
	- 0.20	Guidelines
Frac <sub>leach</sub>	Fraction of N that area lost through leaching and	Table 11.3 of Ch. 11
	runoff; dimensionless, 0-1	Vol. 4 of 2006 IPCC
	- 0.30	Guidelines

# History of the document

|

Version	Date	Contents revised
01.0	21 February 2020	JC5, Annex 5
		Initial approval.